



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which may be denoted by T_0 , another part T_1 , involves the first powers, and a third T_2 involves the squares and products of the same; then $T = T_0 + T_1 + T_2$. By the theory of homogeneous functions

$$\sum_i (p_i q_i') = \sum_i \left(\frac{dT}{dq_i'} q_i' \right) = T_1 + 2T_2.$$

Hence, writing

$$\mathcal{Q}' = \mathcal{Q} + T_0,$$


$$H = T_2 - \mathcal{Q}'.$$

GEOMETRICAL PROBLEM.

BY T. P. STOWELL, ROCHESTER, N. Y.

In any triangle whose angles are all known, as also the distances from the three angles to any point either within or without the triangle, the sides may be found as follows:

Let ABC be the triangle, and P a point whose distances PA, PB and PC are given; revolve the triangle APC over the side AC, P will be at F; and APB revolved over AB, P will be at D; also BPC revolved over BC, P will be at E; then join DE, EF and FD.



The $\triangle AFD$ is known, having two equal sides, AF and AD, and the included angle DAF double the angle BAC. Hence DF is known. Similarly FE and DE are

known, and therefore all the angles of the $\triangle DEF$ are known. Consequently $\angle APC = AFD + DFE + CFE$, and therefore the sides AC, AB and BC are easily found. This question admits of an easy geometrical construction.

NOTE ON THE SOLUTION OF QUADRATIC EQUATIONS, BY WM. ROCK, C. E., EAGLE VILLAGE, N. Y.—The following method of solving Quadratic Equations does not appear in many excellent books on Algebra. The object of the method is to solve any quadratic equation without involving fractions.

$$\text{Let } ax^2 + 2bx = n \dots\dots\dots (1)$$

be any quadratic. Assume $x = y \div a$, then will $ax^2 = y^2 \div a$, and $2bx = 2by \div a$. Substitute these values in the given equation and we have

$$\frac{y^2}{a} + \frac{2by}{a} = n \dots\dots\dots (2).$$

Multiply (2) by a and we have

$$y^2 + 2by = an \dots\dots\dots (3).$$

We have thus removed the coefficient of x^2 without changing that of x . Solving (3) by the ordinary rule, we have

$$y = -b \pm \sqrt{an + b^2}.$$

Hence

$$x = \frac{y}{a} = \frac{-b \pm \sqrt{an + b^2}}{a},$$

which gives the following arithmetical

Rule: If the coefficient of x is an odd number double the whole equation, then multiply the right hand member by the coefficient of x^2 , to the product add the square of half the coefficient of x , extract the square root of the result, connect this square root with half the coefficient of x with its sign changed and divide this result by the coefficient of x^2 ; the quotient is the value of x .

Numerical Example.—Let $11x^2 - 17x = 108$.

Since the coefficient of x is odd, double the equation and apply the rule to the resulting equation:

$22x^2 - 34x = 216$. Then $216 \times 22 = 4752$, add $(\frac{1}{2} \times 34)^2 = 17^2 = 289$, and we have $4752 + 289 = 5041$, extracting square root of this sum we have $\sqrt{5041} = \pm 71$. Hence

$$x = \frac{17 \pm 71}{22} = 4 \text{ or } -2\frac{5}{11}.$$